

On the link between Shrödinger and Vlasov equations

Tigran Aivazian ¹

Abstract

It is shown that well-known Vlasov equation can be derived by adding "hidden" degrees of freedom and subsequent quantization. The Shrödinger equation obtained in this manner coincides (in x-representation) with the kinetic equation for the original dynamical system

1. Two ways to derive the Vlasov equation

Consider a non-autonomous dynamical system in R^n :

$$\frac{dx^i}{dt} = X^i(x^1, \dots, x^n, t) \quad (1.1)$$

We assume that the vector field X is solenoidal:

$$\operatorname{div} X \equiv \frac{\partial X^i}{\partial x^i} = 0 \quad (1.2)$$

One can derive Vlasov equation (known also as Liouville or collisionless Boltzmann) from the condition that the integral of the distribution function $f(x, t)$ is invariant with respect to the phase flow generated by the field X :

$$\int_G f(x, 0) d^n x = \int_{g^t(G)} f(x, t) d^n x, \quad (1.3)$$

where $G \subseteq R^n$ is a open subset in R^n , $g^t(x)$ is a trajectory starting at point $x \in G$ and $d^n x$ is a standard integration measure in Euclidean space. The Vlasov equation has the following form:

$$\frac{\partial f}{\partial t} + \frac{\partial}{\partial x^i} \{X^i f\} = 0 \quad (1.4)$$

Statement. Kinetic equation can be also obtained by doubling the dimension of the base phase space and subsequent quantization of the resulting hamiltonian system. Indeed, one can add n variables p_i and a function $H(x, p, t)$ so that the dynamical system becomes hamiltonian:

$$\begin{aligned} \dot{x}^i &= \frac{\partial H}{\partial p_i} = X^i(x, t), \\ \dot{p}_i &= -\frac{\partial H}{\partial x^i} = -p_k \frac{\partial X^k(x, t)}{\partial x^i}, \\ H(x, p, t) &= p_i X^i(x, t) \end{aligned} \quad (1.5)$$

¹Please use email tigran@zetnet.co.uk for correspondence

Since this system is hamiltonian it is possible to construct the corresponding equations of quantum mechanis for it:

$$i\frac{\partial\psi}{\partial t} = \hat{H}\psi \quad (1.6)$$

Using the condition of divergenless of X we rewrite the last equation as:

$$\frac{\partial\psi}{\partial t} + X^i\frac{\partial\psi}{\partial x^i} = 0, \quad (1.7)$$

As we see the Shrödinger equation for wave function $\psi(x, t)$ in x-representation coincides with the Vlasov equation for distribution function $f(x, t)$ which proves our original statement.